


Criteria for Selection of Patients for Same-Day Discharge after PCI

Felton Hospital¹



1) Pseudonym.

Source: Cardiovascular Roundtable interviews and analysis.

Criteria for Selection of Patients for Same-Day Discharge after PCI (Cont.)

Felton Hospital¹

Pre-Admission Criteria

The following is completed by the Physician/NP/PA in the pre-admission area and follows the patient through to discharge.

MAJOR CRITERIA	
Acute Coronary Syndrome	
Clinical Heart Failure within 7 days	
Creatinine greater than 1.5 mg/dl, GFR less than 60 ml/min, Hemodialysis	
Inability to ambulate due to poor coordination, vasomotor instability, dizziness or suspected neurologic issues or events	
Need for therapeutic anticoagulation (i.e. Coumadin)	
Left Ventricular dysfunction (Ejection Fraction less than 30%)	
Antiplatelet or contrast allergy (uncontrolled)	
Prior stent thrombosis	

MINOR CRITERIA	
Age greater than 75 years	
Patient lives or will be staying greater than 60 minutes from hospital	
Patient does not live with or have the ability to stay with, another responsible adult on the evening of intended discharge	
History of recent ventricular arrhythmia	
Uncontrolled Diabetes (HgA1c > 7% or Fasting Blood Sugar > 180 mg/dl)	
Poorly controlled Hypertension (SBP > 160 mmHg)	
COPD (Home Oxygen dependent or requiring acute treatment)	
Morbidly obese	
Severe valvular heart disease (ie-Severe AS, Severe MR)	
Prior organ transplant/immunosuppressive therapy	

1) Pseudonym.

Source: Cardiovascular Roundtable interviews and analysis.

Criteria for Selection of Patients for Same-Day Discharge after PCI (Cont.)

Felton Hospital¹

Procedure Criteria

(To be completed by the Interventional Cardiologist)

MAJOR	
Complex Anatomy: Unprotected Left Main, PCI performed in 2 or more vessels, Bifurcation Lesion or Long Lesion (≥ 28 mm), Support device (i.e. Balloon Pump, Impella)	
Suboptimal angiographic results (i.e. residual dissection)	
Use of GPIIb-IIIa Inhibitors (i.e. Abciximab, Eptifibatide)	
Side branch occlusion	
No-Reflow requiring treatment	
Significant Hypertension or Hypotension during procedure (SBP < 100 mmHg or > 160 mmHg)	
Total contrast volume/Calculated CrCl (CCC) ratio > 3	
Intra-procedural access site complication	
Significant brady or tachy arrhythmia requiring treatment	

MINOR	
Rotablator, Laser	
Chronic Total Occlusion	
Proximal LAD lesion	
SVG PCI	
Fluoro > 60 minutes or > 5 Gy	
Diffuse in-stent restenosis with restenting	
Left Ventricular End Diastolic Pressure > 25 mmHg	

1) Pseudonym.

Source: Cardiovascular Roundtable interviews and analysis.

Criteria for Selection of Patients for Same-Day Discharge after PCI (Cont.)

Felton Hospital¹

Criteria for Admission

INPATIENT Admission

(1 **MAJOR** criteria / 2 or more **MINOR** criteria)
Patient required an inpatient admission due to their co-morbid conditions and complex/high risk nature of their procedure and was considered to be at increased risk of major adverse cardiac and vascular events if discharged.

Ambulatory Encounter:

- SAME DAY DISCHARGE**
 EXTENDED RECOVERY (Patient requires overnight recovery)

Physician Signature:

Date ___ / ___ / ___
Time ___:___ AM / PM

1) Pseudonym.

Source: Cardiovascular Roundtable interviews and analysis.